## Indexes

## **Author Index**

Abdul-Rahim, H. see Husseini, A.

Abouaesha, F. see Ovibo, S.O.

Abraha, H.D. see Mcintyre, E.A.

Abrahamsen, B., Rohold, A., Henriksen, J.E. & Beck-Nielsen, H.

Correlations between insulin sensitivity and bone mineral
density in non-diabetic men, 124

Abrams, C. see Meadows, K.A.

Agarwal, M.M., Hughes, P.F, Punnose, J. & Ezimokhai, M. Fasting plasma glucose as a screening test for gestational diabetes in a multi-racial, high-risk population, 720

Ahluwalia, A., Sood, A., Sood, A., Lakshmy, R., Kapil, A. & Pandey, R.M. Nasal colonization with *Staphylococcus aureus* in patients with diabetes mellitus, 487

Airey, C.M., Williams, D.D.R., Martin, P.G., Bennett, C.M.T. & Spoor, P.A. Hypoglycaemia induced by exogenous insulin – 'human' and animal insulin compared, 416

Åkerblom, H.K. see Tupola, S.

Allard, S. see Meeking, D.R.

Antonellis, A. see Moczulski, D.K.

Arnqvist, H. see Öestman, J.

Ashkenazi, I. see Kida, K.

Avery, L. see Browne, D.L.

Awartani, F. see Husseini, A.

Bache, I. see Kyvik, K.O.

Bailey, C.J. The rise and fall of troglitazone, 414

Baker, N.R. see Yuen, K.C.J.

Baker, R. see Hutchinson, A.

Balent, B. see Brunner, G.A.

Bardini, G. see Mannucci, E.

Barnett, A.H. see Lloyd, C.E.

Barnett, A.H. see Mcternan, C.L.

Baroni, A. see Frontoni, S.

Barrett, P. see Mckenna, K.

Barrett, T.G. see Ehtisham, S.

Barth, J.H. see Marshall, S.M.

Barth, J.H. see Thomas, A.

Bartlett, W.A., Jones, A.F., Hall, R.A. & Legg, E.F. Standardization of HbA<sub>1c</sub> measurements, 558

Basciani, F. see Chiarelli, F.

Becker, M. see Kordonouri, O.

Beck-Nielsen, H. see Abrahamsen, B.

Beck-Nielsen, H. see Fenger, M.

Beck-Nielsen, H. see Jensen, D.M.

Beck-Nielsen, H. see Kyvik, K.O.

Beeching, N. see Gill, G.

Beeching, N.J. see Gill, G.

Begley, J.P. & Kerr, D. Diábetes and lipids – a conundrum,

Bender, R. see Mühlhauser, I.

Bennett, C.M.T. see Airey, C.M.

Bennett, P.H. Type 2 Diabetes: Prediction and Prevention, 625

Berger, M. see Bott, U.

Berger, M. see Mühlhauser, I.

Berglund, G. see Hedblad, B.

Best, C. see Coppini, D.V.

Bhatia, E. see Singh, A.K.

Bhatia, V. see Singh, A.K.

Bilous, R.W. see New, J.P.

Bingham, C., Nicholls, A.J. & Hattersley, A.T. Heterogeneity in the clinical course of patients with Type 2 diabetes on dialysis the need for different preventative strategies, 685

Biertness, E. see Husseini, A.

Björk, E. see Öestman, J.

Blohmé, G. see Öestman, J.

Bo, S., Cavallo-Perin, P., Gentile, L., Repetti, E. & Pagano, G. Influence of a familial history of diabetes on the clinical characteristics of patients with Type 2 diabetes mellitus, 538

Bo, S., Cavallo-Perin, P., Scaglione, L., Ciccione, G. & Pagano, G. Low birthweight and metabolic abnormalities in twins with increased susceptibility to Type 2 diabetes mellitus, 365

Bodansky, H.J. see Fear, N.T.

Bodansky, H.J. see Law, G.R.

Bodansky, H.J. see Mckinney, P.A.

Bolinder, J. see Öestman, J.

Bonnaud, S. see Rigalleau, V.

Booth, A. see Hutchinson, A.

Borch-Johnsen, K. see Wareham, N.J.

Borg, K. see Johansson, B.-L.

Bott, S. see Bott, U.

Bott, U., Bott, S., Hemmann, D. & Berger, M. Evaluation of a holistic treatment and teaching programme for patients with Type 1 diabetes who failed to achieve their therapeutic goals under intensified insulin therapy, 635

Boulton, A.J.M. see Khawaja, K.I.

Boulton, A.J.M. see Oyibo, S.O.

Bourdel-Marchasson, I., Dubroca, B., Letenneur, L., Fourrier, A., Richard-Harston, S., Decamps, A., Rainfray, M., Emeriau, J.-P. & Dartigues, J.-F. Incidence and predictors of drug-treated diabetes in elderly french subjects. The PAQUID Epidemiological Survey, 675

Bourgeois, M.L. see Rigalleau, V.

Bracaglia, D. see Frontoni, S.

Bradshaw, D. see Levitt, N.S.

Brady, S. see Rutter, M.K.

Browne, B.H. see Gordon, D.

Browne, D.L., Avery, L., Turner, B.C., Kerr, D. & Cavan, D.A.
What do patients with diabetes know about their tablets?,

Bruce, D.G. see Davis, T.M.E.

Brunner, G.A., Hirschberger, S., Sendlhofer, G., Wutte, A., Ellmerer, M., Balent, B., Schaupp, L., Krejs, G.J. & Pieber, T.R. Post-prandial administration of the insulin analogue insulin aspart in patients with Type 1 diabetes mellitus, 371

Burden, A.C. see Davies, M.J.

Buschard, K. see Kyvik, K.O.

Buter, H., Van Tol, A., Navis, G.J., Scheek, L.M., De Jong, P.E., De Zeeuw, D. & Dullaart, R.P.F. Angiotensin-II receptor antagonist treatment lowers plasma total and very low+low-density lipoprotein cholesterol in Type 1 diabetic patients with albuminuria without affecting plasma cholesterol esterification and cholesteryl ester transfer, 550

Butler, C.C. see Morgan, C.Ll.

Buyken, A.E., Toeller, M., Heitkamp, G., Irsigler, K., Holler, C., Santeusanio, F., Stehle, P., Fuller, J.H. & Eurodiab Iddm Complications Study Group, J.H. Carbohydrate sources and glycaemic control in Type 1 diabetes mellitus, 351

Cameron, F. see Wake, M.

Campbell, I.W. see Song, S.H.

Campbell, L.V. see Depczynski, B.

Cappoen, J.P. see Vambergue, A.

Caputo, S. see Ghirlanda, G.

Carlsson, S., Hammar, N., Efendic, S., Persson, P.-G., Östenson, C.-G. & Grill, V. Alcohol consumption, Type 2 diabetes mellitus and impaired glucose tolerance in middle-aged Swedish men, 776

Carmona, F. see Costa, A.

Carney, T.A., Guy, S.P. & Helliwell, C.D. Seasonal variation in  $HbA_{1c}$  in patients with Type 2 diabetes mellitus, 554

Caselli, A. see Frontoni, S.

Catalan, J. see Mitchell, C.

Cavallo-Perin, P. see Bo, S.

Cavan, D. see Flanagan, D.E.H.

Cavan, D.A. see Browne, D.L.

Chan, J.C.N. see Li, J.K.Y.

Cheetham, C. see Collis, I.

Chen, C.-Y. see Hsu, T.-C.

Chew, S.K. see Tai, E.S.

Chiarelli, F., Spagnoli, A., Basciani, F., Tumini, S., Mezzetti, A., Cipollone, F., Cuccurullo, F., Morgese, G. & Verrotti, A. Vascular endothelial growth factor (VEGF) in children, adolescents and young adults with Type 1 diabetes mellitus: relation to glycaemic control and microvascular complications, 650

Chiou, J.Y.'see Jap, T.-S.

Chisholm, D.J. see Depczynski, B.

Chisholm, D.S. see Depczynski, B.

Chowienczyk, P.J. see Meeking, D.R.

Ciani, S. see Mannucci, E.

Ciccione, G. see Bo, S.

Cinek, O., Lánská, V., Kolousková, S., Šumník, Z., Snajderová, M., Rønningen, K.S., Vavrinec, J. & Collaborators Of The Czech Childhood Diabetes Registry, J. Type 1 diabetes mellitus in Czech children diagnosed between 1990–1997: a significant increase in incidence and male predominance in the age group 0–4 years, 64

Cipollone, F. see Chiarelli, F.

Clark, P.M. see Whitelaw, D.C.

Clausen, P., Jacobsen, P., Rossing, K., Jensen, J.S., Parving, H.H. & Feldt-Rasmussen, B. Plasma concentrations of VCAM-1 and ICAM-1 are elevated in patients with Type 1 diabetes mellitus with microalbuminuria and overt nephropathy, 644

Clowes, J. see Garvican, L.

Cockram, C.S. see Janus, E.D.

Cockram, C.S. see Li, I.K.Y.

Coffey, M. see Griffin, M.E.

Collins, P.B. see Owens, D.

Collis, J., Cheetham, C., Dembo, L., O'Driscoll, J., Stanton, K., Taylor, R. & Green, D. Losartan, an angiotensin type 1 receptor inhibitor, and endothelial vasodilator function in Type 1 diabetes mellitus, 553

Colman, P.G. see Singh, A.K.

Conget, I. see Costa, A.

Connell, J. see Mckenna, K.

Connor, H. see Nelson, M.

Connor, H. see Oyibo, S.O.

Conroy, C. see Simmons, D.

Coppini, D.V. & Best, C. A case of hand ulceration in the diabetic foot clinic – a reminder of hand neuropathy in 'at risk' patients,

Coppini, D.V. see Weng, C.

Coppini, D.V., Weng, C., Young, P.J. & Sönksen, P.H. The 'VPTscore' – a useful predictor of neuropathy in diabetic patients. 488

Corrall, R.I. see Pearson, E.R.

Costa Forti, A. see Schmidt, M.I.

Costa, A., Carmona, F., Martinez-Roman, S., Quintó, L., Levy, I. & Conget, I. Post-partum reclassification of glucose tolerance in women previously diagnosed with gestational diabetes mellitus. 595

Coster, S., Gulliford, M.C., Seed, P.T., Powrie, J.K. & Swaminathan, R. Self-monitoring in Type 2 diabetes mellitus: a meta-analysis. 755

Cotroneo, P. see Ghirlanda, G.

Crace, C.J. Exercise and Sport in Diabetes, 333

Cremasco, F. see Mannucci, E.

Cuccurullo, F. see Chiarelli, F.

Cummings, E.A. see O'Hayon, B.E.

Cummings, M.H. see Meeking, D.R.

Cundy, T., Gamble, G., Townend, K., Henley, P.G., Macpherson, P. & Roberts, A.B. Perinatal mortality in Type 2 diabetes mellitus, 33

Currie, C.J. see Dixon, S.

Currie, C.J. see Morgan, C.Ll.

Dabadghao, P. see Singh, A.K.

Dahlin, L.-B. see Sundkvist, G.

Dalton, R.N. see Schultz, C.J.

Daly, B. see Depczynski, B.

Damm, P. see Hellmuth, E.

Daneman, D. see O'Havon, B.E.

Danne, T. see Kordonouri, O.

Dartigues, J.-F. see Bourdel-Marchasson, I.

Davies, C. see Rajbhandari, S.

Davies, M.J., Raymond, N.T., Day, J.L., Hales, C.N. & Burden, A.C. Impaired glucose tolerance and fasting hyperglycaemia have different characteristics. 433

Davis, T.M.E., Zimmet, P., Davis, W.A., Bruce, D.G., Fida, S. & Mackay, I.R. Autoantibodies to glutamic acid decarboxylase in diabetic patients from a multi-ethnic Australian community: the Fremantle Diabetes Study, 667

Davis, W.A. see Davis, T.M.E.

Day, J.L. see Davies, M.J.

De Angelis, C. see Frontoni, S.

De Block, C.E.M., De Leeuw, I.H., Rooman, R.P.A., Winnock, F.,
Du Caju, M.V.L., Van Gaal, L.F. & The Belgian Diabetes
Registry. Gastric parietal cell antibodies are associated with
glutamic acid decarboxylase-65 antibodies and the HLA
DQA1\*0501-DQB1\*0301 haplotype in Type 1 diabetes
mellitus, 618

De Jong, P.E. see Buter, H.

De Leeuw, I.H. see De Block, C.E.M.

De Lima, L. see Schmidt, M.I.

De Zeeuw, D. see Buter, H.

Deacon, L.F. see Nolan, J.J.

Decamps, A. see Bourdel-Marchasson, I.

Deepa, R., Shanthi, S., Rani, G., Premalatha, V. & Mohan, V. Comparison of ADA 1997 and WHO 1985 criteria for diabetes in south Indians – the Chennai Urban Population Study, 872

Dekker, J.H. see Franse, L.V.

Demaine, A.G. see Zhao, H.X.

Dembo, L. see Collis, J.

Demuth, H.-U. see Meneilly, G.S.

Depczynski, B., Daly, B., Campbell, L.V., Chisholm, D.J. & Keogh, A. Predicting the occurrence of diabetes mellitus in recipients of heart transplants. 15

Depczynski, B., Daly, B., Campbell, L.V., Chisholm, D.S. & Keogh, A. Authors' response to the letter 'Predicting the occurrence of diabetes mellitus in recipients of heart transplants', 624

Dieterich, W. see Kordonouri, O.

Dixon, S., Currie, C.J. & Peters, J.R. The cost of diabeties: time for a different approach?, 820

Dognin, C. see Vambergue, A.

Dogra, G. & Watts, G.F. Endothelial dysfunction in Type 1 diabetes mellitus: methodological considerations, 687

Donnan, P. see Evans, J.

Donohoe, M.E., Felton, J.A., Hook, A., Powell, R., Robinson, I., Stead, J.W., Taylor, R. & Tooke, J.E. Improving foot care for people with diabetes mellitus – a randomized controlled trial of an integrated care approach, 581

Dorling, M.W. see Petry, C.J.

Dornan, T. Diabetes: Emergency and Hospital Management, 816

Dornhorst, A. see Frost, G.

Dornhorst, A. see Packer, S.C.

Du Caju, M.V.L. see De Block, C.E.M.

Dubroca, B. see Bourdel-Marchasson, I.

Dullaart, R.P.F. see Buter, H.

Dunger, D. see Schultz, C.J.

Dunne, M.J. Ions, genes and insulin release: from basic science to clinical disease, 91

Dver, P.H. see Lloyd, C.E.

Edelsten, C. see Yuen, K.C.J.

Edmonds, M. see Winkler, A.S.

Efendic, S. see Carlsson, S.

Ehtisham, S., Barrett, T.G. & Shaw, N.J. Type 2 diabetes mellitus in UK children – an emerging problem, 867

Eiskiaer, N. see Winkler, A.S.

Elliott, T. see Meneilly, G.S.

Ellis, I. see Evans, I.

Ellmerer, M. see Brunner, G.A.

Emeriau, J.-P. see Bourdel-Marchasson, I.

Endo, G. see Okada, K.

Ericsson, M. see Reichard, P.

Eriksson, J. see Öestman, J.

Eriksson, K.-F. see Sundkvist, G.

Eurodiab IDDM Complications Study Group. see Buyken, A.E.

Evans, J., Ellis, J., Steinke, D., Donnan, P. & Morris, A.D. Childhood Type 1 diabetes and parental occupation involving

social mixing and infectious contacts, 490 Evans, J.D. see Morgan, C.Ll.

Evans, J.M.M., Newton, R.W., Ruta, D.A., Macdonald, T.M. & Morris, A.D. Socio-economic status, obesity and prevalence of Type 1 and Type 2 diabetes mellitus, 478

Everett, J. see Flanagan, D.E.H.

Ezimokhai, M. see Agarwal, M.M.

Fanelli, A. see Mannucci, E.

Farrace, S. see Frontoni, S.

Fear, N.T., Mckinney, P.A., Parslow, R.C., Patterson, C.C. & Bodansky, H.J. Childhood Type 1 diabetes and parental occupation involving social mixing and infectious contacts. Reply. 491

Feher, M.D. see Mitchell, C.

Feher, M.D. see Munro, N.

Feilberg-Jørgensen, N. see Jensen, D.M.

Feldt-Rasmussen, B. see Clausen, P.

Felton, J.A. see Donohoe, M.E.

Fenger, M., Poulsen, P., Beck-Nielsen, H. & Vaag, A. Impact of the Xba1-polymorphism of the human muscle glycogen synthase gene on parameters of the insulin resistance syndrome in a Danish twin population, 735

Fernqvist-Forbes, E. see Johansson, B.-L.

Fida, S. see Davis, T.M.E.

Finer, M.L. see Reviewed By, N.

Firth, R.G. see Griffin, M.E.

Flanagan, D.E.H., Watson, J., Everett, J., Cavan, D. & Kerr, D. Driving and insulin – consensus, conflict or confusion?, 316

Fogarty, L. see Vaughan, N.J.A.

Foley, M. see Griffin, M.E.

Fontaine, P. see Vambergue, A.

Forrester, J.V. see Hipwell, J.H.

Forster, J. see Rutter, M.K.

Fourrier, A. see Bourdel-Marchasson, I.

Franse, L.V., Valk, G.D., Dekker, J.H., Heine, R.J. & Van Eijk, J.T.M. Numbness of the feet' is a poor indicator for polyneuropathy in Type 2 diabetic patients, 105 Frier, B.M. see Perros, P.

Frier, B.M. see Strachan, M.W.J.

Frontoni, S., Pellegrinotti, M., Bracaglia, D., Farrace, S., Caselli, A., Baroni, A., De Angelis, C., Menzinger, G. & Gambardella, S. Hyperinsulinaemia in offspring of Type 2 diabetic patients: impaired response of carbohydrate metabolism, but preserved cardiovascular response, 606

Frost, G. & Dornhorst, A. The relevance of the glycaemic index to our understanding of dietary carbohydrates, 336

Frost, G.S. see Packer, S.C.

Fuecker, K. see Hanefeld, M.

Fuiii, S. see Okada, K.

Fuller, J.H. see Buyken, A.E.

Gale, E.A.M. For The UK Trial Group. A randomized, controlled trial comparing insulin lispro with human soluble insulin in patients with Type 1 diabetes on intensified insulin therapy, 209

Gallacher, B. see Meier, M.

Gambardella, S. see Frontoni, S.

Gamble, G. see Cundy, T.

Garvican, L., Clowes, J. & Gillow, T. Preservation of sight in diabetes: developing a national risk reduction programme, 627

Gatta, B. see Rigalleau, V.

Gellert, S.A. see Singh, A.K.

Gentile, L. see Bo, S.

George, E. see Sturrock, N.D.C.

Giacaman, R. see Husseini, A.

Gibbins, R.L. see Owens, D.R.

Gill, G. & Beeching, N. Capturing the prevalence of diagnosed diabetes, 753

Gill, G. & Beeching, N.J. Deprivation and Type 2 diabetes mellitus prevalence, 813

Gill, G.V. see Ismail, A.A.

Gillow, T. see Garvican, L.

Gin, H. see Rigalleau, V.

Glen, A. see Mckenna, K.

Glenn, A. see Mckenna, K.

Gold, A. Hypoglycaemia in Clinical Diabetes, 411

Gomis, R. see Wolffenbuttel, B.H.R.

Gordon, D. & Browne, B.H. Macular oedema with associated uveitis and cataract following presentation of Type 1 diabetes mellitus in severe ketoacidosis, 326

Goyder, E.C. & Irwig, L.M. Screening for Type 2 diabetes mellitus: a decision analytic approach, 469

Green, A. see Kyvik, K.O.

Green, D. see Collis, I.

Greene, D.A. see Sundkvist, G.

Greenwood, R. see Owens, D.R.

Griffin, M.E., Coffey, M., Johnson, H., Scanlon, P., Foley, M., Stronge, J., O'Meara, N.M. & Firth, R.G. Universal vs. risk factor-based screening for gestational diabetes mellitus: detection rates, gestation at diagnosis and outcome, 26

Grill, V. see Carlsson, S.

Grimshaw, G.M. see Owens, D.R.

Groop, L. see Klannemark, M.

Gubina, M. see Urbancic-Rovan, V.

Gulliford, M.C. see Coster, S.

Gurney, K.A. see McKinney, P.A. Guy, S.P. see Carney, T.A.

Hales, C.N. see Davies, M.J.

Hall, R.A. see Bartlett, W.A.

Hammar, N. see Carlsson, S.

Hanefeld, M., Koehler, C., Henkel, E., Fuecker, K., Schaper, F. & Temelkova-Kurktschiev, T. Post-challenge hyperglycaemia relates more strongly than fasting hyperglycaemia with carotid intima-media thickness: the RIAD Study, 835

Harding, S. see Owens, D.R.

Hart, L.M.'t see Van Essen, E.H.R.

Hartland, B. see Nelson, M.

Hatterslev, A.T. see Bingham, C.

Hattersley, A.T. see Pearson, E.R.

Hayashi, T. see Okada, K.

Hayat, S.A. see Khawaja, K.I.

Hedblad, B., Nilsson, P., Janzon, L. & Berglund, G. Relation between insulin resistance and carotid intima-media thickness and stenosis in non-diabetic subjects. Results from a crosssectional study in Malmö, Sweden, 299

Hedley, A.J. see Lam, T.H.

Heine, R.J. see Franse, L.V.

Heitkamp, G. see Buyken, A.E.

Heller, S. see Morris, B.W.

Heller, S.R. Reflections of a new Editor, 819

Heller, S.R. New clothes for Diabetic Medicine, 413

Heller, S.R. The last editor of Diabetic Medicine?, 1

Heller, S.R. The quality of clinical care in diabetes in Britain – could do better, 335

Helliwell, C.D. see Carney, T.A.

Hellmuth, E., Damm, P. & Mølsted-Pedersen, L. Oral hypoglycaemic agents in 118 diabetic pregnancies, 507

Hemmann, D. see Bott, U.

Heng, D. see Tai, E.S.

Henkel, E. see Hanefeld, M.

Henley, P.G. see Cundy, T.

Henriksen, J.E. see Abrahamsen, B.

Hesketh, K. see Wake, M.

Hipwell, J.H., Strachan, F., Olson, J.A., Mchardy, K.C., Sharp, P.F. & Forrester, J.V. Automated detection of microaneurysms in digital red-free photographs: a diabetic retinopathy screening tool, 588

Hirschberger, S. see Brunner, G.A.

Holler, C. see Buyken, A.E.

Home, P.D., Lindholm, A. & Riis For The European Insulin Aspart Study Group, A. Insulin aspart vs. human insulin in the management of long-term blood glucose control in Type 1 diabetes mellitus: a randomized controlled trial, 762

Hook, A. see Donohoe, M.E.

Houghton, G.M. see Ismail, A.A.

Hsu, T.-C., Wang, C.-L., Shau, Y.-W., Tang, F.-T., Li, K.-L. & Chen, C.-Y. Altered heel-pad mechanical properties in patients with Type 2 diabetes mellitus, 854

Hughes, P.F see Agarwal, M.M.

Hunt, J. see Morgan, C.Ll.

Husseini, A., Abdul-Rahim, H., Awartani, F., Giacaman, R., Jervell, J. & Bjertness, E. Type 2 diabetes mellitus, impaired glucose tolerance, and associated factors in a rural Palestinian village, 746

Hutchinson, A., Mcintosh, A., Peters, J., O'Keeffe, C., Khunti, K., Baker, R. & Booth, A. Effectiveness of screening and monitoring tests for diabetic retinopathy – a systematic review, 495

Huttunen, N.P. see Tupola, S.

Irsigler, K. see Buyken, A.E.

Irwig, L.M. see Goyder, E.C.

Ismail, A.A., Gill, G.V., Lawton, K., Houghton, G.M. & Macfarlane, I.A. Comparison of questionnaire, breath carbon monoxide and urine cotinine in assessing the smoking habits of Type 2 diabetic patients, 119

Ismail, A.A., Wallymahmed, M.E., Gill, G.V. & MacFarlane, I.A. Failure to reduce nicotine addiction in young adults with diabetes, 330

Ito, T. see Kida, K.

Jacobsen, P. see Clausen, P.

Jansen, J.J. see van Essen, E.H.R.

Janus, E.D. see Lam, T.H.

Janus, E.D., Wat, N.M.S., Lam, K.S.L., Cockram, C.S., Siu, S.T.S., Liu, L.J. & Lam, T.H. On Behalf Of The Hong Kong Cardiovascular Risk Factor Steering Committee. The prevalence of diabetes, association with cardiovascular risk factors and implications of diagnostic criteria (ADA 1997 and WHO 1998) in a 1996 community-based population study in Hong Kong Chinese, 741

Janzon, L. see Hedblad, B.

Jap, T.-S., Wu, Y.-C., Chiou, J.Y. & Kwok, C.-F. A novel mutation in the hepatocyte nuclear factor-1α/MODY3 gene among Chinese subjects with early-onset Type 2 diabetes mellitus in Taiwan, 390

Jeffcoate, W. The Care of Neuropathic Limbs: A Practical Manual, 169

Jeffcoate, W., Lima, J. & Nobrega, L. The Charcot foot, 253
Jensen, D.M., Sørensen, B., Feilberg-Jørgensen, N., Westergaard,
J.G. & Beck-Nielsen, H. Maternal and perinatal outcomes in
143 Danish women with gestational diabetes mellitus and 143
controls with a similar risk profile, 281

Jensen, J.S. see Clausen, P.

Jensen-Urstad, K. see Reichard, P.

Jensen-Urstad, M. see Reichard, P.

Jervell, J. see Husseini, A.

Joffe, B.I. see Panz, V.R.

Johansson, B.-L., Borg, K., Fernqvist-Forbes, E., Kernell, A., Odergren, T. & Wahren, J. Beneficial effects of C-peptide on incipient nephropathy and neuropathy in patients with Type 1 diabetes mellitus, 181

John, G. see Thomas, A.

Johnson, A. see Owens, D.

Johnson, H. see Griffin, M.E.

Johnston, D.G. see Kousta, E.

Jones, A.F. see Bartlett, W.A.

Jones, N.P. see Nolan, J.J.

Jones, N.P. see Wolffenbuttel, B.H.R.

Jorde, R. & Sundsfjord, J. Intra-individual variability and longitudinal changes in glycaemic control in patients with Type 1 diabetes mellitus, 451 Jörgens, V. see Mühlhauser, I.

Joubert, G. see Levitt, N.S.

Juneja, R. see Meneilly, G.S.

Kalk, W.J. see Panz, V.R.

Kapil, A. see Ahluwalia, A.

Keogh, A. see Depczynski, B.

Kernell, A. see Johansson, B.-L.

Kerr, D. see Beglev, J.P.

Kerr, D. see Browne, D.L.

Kerr, D. see Flanagan, D.E.H.

Kesson, C.M. see Mckenna, K.

Khawaja, K.I., Walker, D., Hayat, S.A., Boulton, A.J.M. & Malik, R.A. Clinico-pathological features of postural hypotension in diabetic autonomic neuropathy, 163

Khunti, K. see Hutchinson, A.

Kida, K., Mimura, G., Ito, T., Murakami, K., Ashkenazi, I., Laron, Z. & The Data Committee For Childhood Diabetes Of The Japan Diabetes Society (JDS). Incidence of Type 1 diabetes mellitus in children aged 0 to 14 in Japan, 1986–1990, including an analysis for seasonality of onset and month of birth: JDS study, 59

Kitange, H.M. see Levitt, N.S.

Kitchen, M.M. see Kong, N.

Klannemark, M., Suurinkeroinen, L., Orho-Melander, M., Groop, L. & Taskinen, M.R. Interaction between the Asn291Ser variant of the *LPL* gene and insulin resistance on dyslipidaemia in high risk individuals for Type 2 diabetes mellitus, 599

Kobayashi, K.-I. see Nagai, Y.

Kobayashi, T. see Oki, Y.

Koehler, C. see Hanefeld, M.

Kohner, E. see Owens, D.R.

Koivukangas, V., Oikarinen, A., Salmela, P.I. & Lahti, A. Microcirculatory response of skin to benzoic acid and methyl nicotinate in patients with diabetes, 130

Kolousková, S. see Cinek, O.

Kong, N., Kitchen, M.M. & Ryder, R.E.J. The use of lispro for high sugar content snacks between meals in intensive insulin regimens, 331

Kordonouri, O., Dieterich, W., Schuppan, D., Weber, G., Müller, C., Sarioglu, N., Becker, M. & Danne, T. Autoantibodies to tissue transglutaminase are sensitive serological parameters for detecting silent coeliac disease in patients with Type 1 diabetes mellitus, 441

Kousta, E., Lawrence, N.J., Penny, A., Millauer, B.A., Robinson, S., Johnston, D.G. & McCarthy, M.I. Women with a history of gestational diabetes of European and South Asian origin are shorter than women with normal glucose tolerance in pregnancy, 792

Kouta, S. see Morris, B.W.

Krejs, G.J. see Brunner, G.A.

Krentz, A.J. Predicting the occurrence of diabetes mellitus in recipients of heart transplants, 488

Krolewski, A.S. see Moczulski, D.K.

Kwok, C.-F. see Jap, T.-S.

Kyvik, K.O., Bache, I., Green, A., Beck-Nielsen, H. & Buschard, K. No association between birth weight and Type 1 diabetes mellitus – a twin-control study, 158 Lahti, A. see Koivukangas, V.

Lakshmy, R. see Ahluwalia, A.

Lam, T.H. see Janus, E.D.

Lam, K.S.L. see Janus, E.D.

Lam, K.S.L. see Lam, T.H.

Lam, T.H., Liu, L.J., Janus, E.D., Lam, K.S.L. & Hedley, A.J. For The Hong Kong Cardiovascular Risk Factor Prevalence Study Steering Committee. Fibrinogen, other cardiovascular risk factors and diabetes mellitus in Hong Kong: a community with high prevalence of Type 2 diabetes mellitus and impaired glucose tolerance, 798

Landin-Olsson, M. see Öestman, J.

Lánská, V. see Cinek, O.

Lao, T.T. see Leung, T.W.

Laron, Z. see Kida, K.

Lattimer, S.A. see Sundkvist, G.

Law, G.R. see McKinney, P.A.

Law, G.R., Bodansky, H.J. & McKinney, P.A. When do environmental factors initiate islet cell damage in children with Type 1 diabetes mellitus?, 167

Lawrence, N.I. see Kousta, E.

Lawson, M.L. see O'Hayon, B.E.

Lawton, K. see Ismail, A.A.

Lean, M.E.I. see Nelson, M.

Legg, E.F. see Bartlett, W.A.

Lemkes, H.H.P.J. see van Essen, E.H.R.

Lernmark, Å. see Öestman, J.

Letenneur, L. see Bourdel-Marchasson, I.

Leung, T.W. & Lao, T.T. Placental size and large-for-gestationalage infants in women with abnormal glucose tolerance in pregnancy, 48

Levitt, N.S., Unwin, N.C., Bradshaw, D., Kitange, H.M., Mbanya, J.-C.N., Mollentze, W.F., Omar, M.A.K., Motala, A.A., Joubert, G., Masuki, G. & Machibya, H. Application of the new ADA criteria for the diagnosis of diabetes to population studies in sub-Saharan Africa, 381

Levy, I. see Costa, A.

Li, J.K.Y., Chan, J.C.N., Zimmet, P.Z., Rowley, M.J., Mackay, I.R. & Cockram, C.S. Young Chinese adults with new onset of diabetic ketoacidosis – clinical course, autoimmune status and progression of pancreatic β-cell function, 295

Li, K.-L. see Hsu, T.-C.

Liddell, W.G. see Pearson, E.R.

Lim, S.C. see Tai, E.S.

Lima, J. see Jeffcoate, W.

Lindblad, L.E. see Reichard, P.

Lindblad, U. see Östgren, C.I.

Lindgärde, F. see Sundkvist, G.

Lindholm, A. see Home, P.D.

Lindström, J., Tuomilehto, J. & Spengler, M. For The Finnish Acarbose Study Group. Acarbose treatment does not change the habitual diet of patients with Type 2 diabetes mellitus, 20

Littorin, B. see Öestman, J.

Liu, L.J. see Janus, E.D.

Liu, L.J. see Lam, T.H.

Lloyd, C.E., Dyer, P.H. & Barnett, A.H. Prevalence of symptoms of depression and anxiety in a diabetes clinic population,

Lord, K. see Nelson, M. Ludlam, C.A. see Perros, P.

Maassen, J.A. see van Essen, E.H.R.

Maccuish, A.C. see Strachan, M.W.J.

MacDonald, T.M. see Evans, J.M.M.

Macarlane, I.A. see Ismail, A.A.

Machibya, H. see Levitt, N.S.

Mackay, I.R. see Davis, T.M.E.

Mackay, I.R. see Li, J.K.Y.

Mackie, A. Handbook of Diabetes (2nd edition), 561

MacNeil, S. see Morris, B.W.

Macpherson, P. see Cundy, T.

Malik, R.A. see Khawaja, K.I.

Mannucci, E., Ognibene, A., Cremasco, F., Bardini, G., Mencucci, A., Pierazzuoli, E., Ciani, S., Fanelli, A., Messeri, G. & Rotella, C.M. Glucagon-like peptide (GLP)-1 and leptin concentrations in obese patients with Type 2 diabetes mellitus, 713

Manto, A. see Ghirlanda, G.

Marshall, S.M. & Barth, J.H. Standardization of HbA<sub>1c</sub> measurements – a consensus statement, 5

Marshall, S.M. see Rutter, M.K.

Marshall, S.M. see Thomas, A.

Martin, P.G. see Airey, C.M.

Martinez-Roman, S. see Costa. A.

Masson, E.A. see Mitchell, L.

Masson, M. see Rigalleau, V.

Masuki, G. see Levitt, N.S. Matos, M.C. see Schmidt, M.I.

Matz, R. Severe diabetic ketoacidosis, 329

Mbanya, J.-C.N. see Levitt, N.S.

McCarthy, M.I. see Kousta, E.

McComb, I.M. see Rutter, M.K.

McGough, N. see Nelson, M.

McGrath, N. Diabetes in New Zealand Maori, 557

McGrath, N.M. & Yeong, M.L. Continuing education, 87

McHardy, K.C. see Hipwell, J.H.

McIntosh, A. see Hutchinson, A.

McIntosh, C. see Mitchell, C.

McIntosh, C. see Munro, N.

McIntosh, C.H.S. see Meneilly, G.S.

McIntyre, E.A., Abraha, H.D., Perros, P. & Sherwood, R.A. Serum S-100β protein is a potential biochemical marker for cerebral oedema complicating severe diabetic ketoacidosis,

McKenna, K., Smith, D., Barrett, P., Glenn, A., Kesson, C.M., Connell, J. & Thompson, C.J. Angiotensin-converting enzyme inhibition by quinapril blocks the albuminuric effect of atrial natriuretic peptide in Type 1 diabetes and microalbuminuria, 219

McKenna, K., Smith, D., Moore, K., Glen, A., Kesson, C.M. & Thompson, C.J. Enhanced albuminuric response to atrial natriuretic peptide in normalbuminuric patients with Type 1 diabetes mellitus – a pilot study, 463

McKenna, K., Smith, D., Tormey, W. & Thompson, C.J. Acute hyperglycaemia causes elevation in plasma atrial natriuretic peptide concentrations in Type 1 diabetes mellitus, 512 McKinney, P.A. see Fear, N.T.

McKinney, P.A. see Law, G.R.

McKinney, P.A., Okasha, M., Parslow, R.C., Law, G.R., Gurney, K.A., Williams, R. & Bodansky, H.J. Early social mixing and childhood Type 1 diabetes mellitus: a case–control study in Yorkshire. UK. 236

McTernan, C.L., Stewart, L.C., Mijovic, C.H. & Barnett, A.H. Assessment of the non-HLA-DR-DQ contribution to IDDM1 in British Caucasian families: analysis of LMP7 polymorphisms, 661

Meadows, K.A., Abrams, C. & Sandbaek, A. Adaptation of the Diabetes Health Profile (DHP-1) for use with patients with Type 2 diabetes mellitus: psychometric evaluation and crosscultural comparison, 572

Meeking, D.R., Allard, S., Munday, J., Chowienczyk, P.J., Shaw, K.M. & Cummings, M.H. Comparison of vasodilator effects of substance P in human forearm vessels of normoalbuminuric Type 1 diabetic and non-diabetic subjects, 243

Meier, M., O'Callaghan, C.J., Gallacher, B., Standl, E. & Williams, B. Hypoxia-induced vascular endothelial growth factor expression is potentiated by elevated glucose levels, 248

Melander, A. see Östgren, C.J.

Mencucci, A. see Mannucci, E.

Meneilly, G.S., Demuth, H.-U., Mcintosh, C.H.S. & Pederson, R.A. Effect of ageing and diabetes on glucose-dependent insulinotropic polypeptide and dipeptidyl peptidase IV responses to oral glucose. 346

Meneilly, G.S., Tildesley, H., Elliott, T., Palmer, J.P. & Juneja, R. Significance of GAD positivity in elderly patients with diabetes. 247

Menzinger, G. see Frontoni, S.

Meredith, I.T. see Skyrme-Jones, R.A.P.

Messeri, G. see Mannucci, E.

Mezzetti, A. see Chiarelli, F.

Middle, J. see Thomas, A.

Mijovic, C.H. see Mcternan, C.L.

Millauer, B.A. see Kousta, E.

Millward, B.A. see Stenhouse, E.

Millward, B.A. see Zhao, H.X.

Mimura, G. see Kida, K.

Mitchell, C., Catalan, J., Mcintosh, C. & Feher, M.D. Joint liaison psychiatry–diabetes clinic: a new specialist service, 481

Mitchell, L. & Masson, E.A. Diabetic foot ulcer management: a perspective of primary and secondary care, 168

Moczulski, D.K., Scott, L., Antonellis, A., Rogus, J.J., Rich, S.S., Warram, J.H. & Krolewski, A.S. Aldose reductase gene polymorphisms and susceptibility to diabetic nephropathy in Type 1 diabetes mellitus, 111

Mohan, V. see Deepa, R.

Mollentze, W.F. see Levitt, N.S.

Mølsted-Pedersen, L. see Hellmuth, E.

Moore, K. see Mckenna, K.

Morgan, C.Ll., Currie, C.J., Hunt, J., Evans, J.D., Rogers, C., Stott, N.C.H. & Peters, J.R. Relative activity and referral patterns for diabetes and non-diabetes in general practice, 230

Morgan, C.Ll., Currie, C.J., Stott, N.C.H., Smithers, M., Butler, C.C. & Peters, J.R. Estimating the prevalence of diagnosed diabetes in a health district of Wales: the importance of using

primary and secondary care sources of ascertainment with adjustment for death and migration, 141

Morgan, C.Ll., Currie, C.J., Stott, N.C.H., Smithers, M., Butler, C.C. & Peters, J.R. The prevalence of multiple diabetes-related complications. 146

Morgese, G. see Chiarelli, F.

Morris, A.D. see Evans, I.

Morris, A.D. see Evans, I.M.M.

Morris, B.W., Kouta, S., Robinson, R., Macneil, S. & Heller, S. Chromium supplementation improves insulin resistance in patients with Type 2 diabetes mellitus, 684

Motala, A.A. see Levitt, N.S.

Mowbray, A. see Song, S.H.

Mühlhauser, I. & Berger, M. Evidence-based patient information in diabetes, 823

Mühlhauser, I., Overmann, H., Bender, R., Jörgens, V. & Berger, M. Predictors of mortality and end-stage diabetic complications in patients with Type 1 diabetes mellitus on intensified insulin therapy, 727

Müller, C. see Kordonouri, O.

Munday, J. see Meeking, D.R.

Munro, N., Riche, N., Mcintosh, C. & Feher, M.D. Too hot to handle – an unusual location for an adverse effect of capsaicin, 552

Murakami, K. see Kida, K.

Murase, T. see Oki, Y.

Nagai, Y., Yamashita, H., Takamura, T. & Kobayashi, K.-I. Does acute hyperinsulinaemia change plasma adrenomedullin concentration in healthy men?. 249

Nakanishi, K. see Oki, Y.

Narasimham, D.V.L. see Vijay, V.

Nattrass, M. see Whitelaw, D.C.

Navis, G.J. see Buter, H.

Neil, H.A.W. see Schultz, C.J.

Nelson, M., Lean, M.E.J., Connor, H., Thomas, B.J., Lord, K., Hartland, B., Waldron, S., Mcgough, N., Walker, L., Ryan, A. & Start, K. Survey of dietetic provision for patients with diabetes, 565

New, J.P., Bilous, R.W. & Walker, M. Insulin sensitivity in hypertensive Type 2 diabetic patients after 1 and 19 days' treatment with trandolapril, 134

Newton, R.W. see Evans, J.M.M.

Nicholls, A.J. see Bingham, C.

Nilsson, H. see Sundkvist, G.

Nilsson, P. see Hedblad, B.

Nobrega, L. see Jeffcoate, W.

Nolan, J.J., Jones, N.P., Patwardhan, R. & Deacon, L.F. Rosiglitazone taken once daily provides effective glycaemic control in patients with Type 2 diabetes mellitus, 287

Nuttens, M.C. see Vambergue, A.

Nuuja, A. see Tupola, S.

Nyström, L. see Öestman, J.

O'Callaghan, C.J. see Meier, M. O'Driscoll, J. see Collis, J.

O'Hayon, B.E., Cummings, E.A., Daneman, D., Ossip, M.G., Lawson, M.L. & Sochett, E.B. Does dietary protein intake correlate with markers suggestive of early diabetic nephropathy in children and adolescents with Type 1 diabetes mellitus?, 708

O'Keeffe, C. see Hutchinson, A.

O'Meara, N.M. see Griffin, M.E.

O'Shea, H. see Song, S.H.

Odergren, T. see Johansson, B.-L.

Öestman, J., Landin-Olsson, M., Törn, C., Palmer, J., Lernmark, Å., Arnqvist, H., Björk, E., Bolinder, J., Blohmé, G., Eriksson, J., Littorin, B., Nyström, L., Scherstén, B., Sundkvist, G. & Wibell, L. Ketoacidosis in young adults is not related to the islet antibodies at the diagnosis of Type 1 diabetes mellitus – a nationwide study, 269

Ognibene, A. see Mannucci, E.

Oikarinen, A. see Koivukangas, V.

Okada, K., Hayashi, T., Tsumura, K., Suematsu, C., Endo, G. & Fujii, S. Leisure-time physical activity at weekends and the risk of Type 2 diabetes mellitus in Japanese men: the Osaka Health Survey, 53

Okasha, M. see McKinney, P.A.

Oki, Y., Okubo, M., Tanaka, S., Nakanishi, K., Kobayashi, T. & Murase, T. Diabetes mellitus secondary to glycogen storage disease type III, 810

Okubo, M. see Oki, Y.

Oldenborg, P.-A., Sundqvist, I.-M. & Sehlin, J. Different effects of glucose on extracellular and intracellular respiratory burst response in normal human neutrophils activated with the soluble agonist fMet-Leu-Phe, 532

Olson, J.A. see Hipwell, J.H.

Omar, M.A.K. see Levitt, N.S.

Orho-Melander, M. see Klannemark, M.

Ossip, M.G. see O'Hayon, B.E.

Östenson, C.-G. see Carlsson, S.

Östgren, C.J., Lindblad, U., Ranstam, J., Melander, A. & Råstam, L. Associations between smoking and β-cell function in a non-hypertensive and non-diabetic population, 445

Overmann, H. see Mühlhauser, I.

Owens, D., Collins, P.B., Johnson, A. & Tomkin, G.H.
Lipoproteins and low-dose estradiol replacement therapy in
post-menopausal Type 2 diabetic patients: the effect of
addition of norethisterone acetate, 308

Owens, D.R., Gibbins, R.L., Kohner, E., Grimshaw, G.M., Greenwood, R. & Harding, S. Diabetic retinopathy screening, 493

Oyibo, S.O., Abouaesha, F., Connor, H. & Boulton, A.J.M. The Diabetic Foot 2000, 875

Ozanne, S.E. see Petry, C.J.

Packer, S.C., Dornhorst, A. & Frost, G.S. The glycaemic index of a range of gluten-free foods, 657

Pagano, G. see Bo, S.

Palmer, J. see Öestman, J.

Palmer, J.P. see Meneilly, G.S.

Pandey, R.M. see Ahluwalia, A.

Pandit, R.J. & Taylor, R. Mydriasis and glaucoma: exploding the myth. A systematic review, 693 Panz, V.R., Kalk, W.J., Zouvanis, M. & Joffe, B.I. Distribution of autoantibodies to glutamic acid decarboxylase across the spectrum of diabetes mellitus seen in South Africa, 524

Parslow, R.C. see Fear, N.T.

Parslow, R.C. see Mckinney, P.A.

Parving, H.H. see Clausen, P.

Patterson, C.C. see Fear, N.T.

Patwardhan, R. see Nolan, J.J.

Patwardhan, R.N. see Wolffenbuttel, B.H.R.

Pawlak, D.B. see Petry, C.J.

Pearson, E.R., Liddell, W.G., Shepherd, M., Corrall, R.J. & Hattersley, A.T. Sensitivity to sulphonylureas in patients with hepatic nuclear factor-1α mutations: evidence for pharmacogenetics in diabetes, 543

Peck, G.M. see Sturrock, N.D.C.

Pederson, R.A. see Meneilly, G.S.

Pellegrinotti, M. see Frontoni, S.

Penny, A. see Kousta, E.

Perros, P. & Frier, B.M. Evaluation of an on-call diabetes service in a large teaching hospital, 386

Perros, P. see McIntyre, E.A.

Perros, P., Singh, R.K., Ludlam, C.A. & Frier, B.M. Prevalence of pernicious anaemia in patients with Type 1 diabetes mellitus and autoimmune thyroid disease, 749

Persson, P.-G. see Carlsson, S.

Peters, J. see Hutchinson, A.

Peters, J.R. see Dixon, S.

Peters, J.R. see Morgan, C.Ll.

Petry, C.J., Dorling, M.W., Wang, C.L., Pawlak, D.B. & Ozanne, S.E. Catecholamine levels and receptor expression in low protein rat offspring, 848

Pieber, T.R. see Brunner, G.A.

Pierazzuoli, E. see Mannucci, E.

Pitocco, D. see Ghirlanda, G.

Poulsen, P. see Fenger, M.

Pound, N. see Sturrock, N.D.C.

Powell, R. see Donohoe, M.E.

Powrie, J.K. see Coster, S.

Premalatha, V. see Deepa, R.

Punnose, J. see Agarwal, M.M.

Quintó, L. see Costa, A.

Rainfray, M. see Bourdel-Marchasson, I.

Rajbhandari, S., Sutton, M., Davies, C., Tesfaye, S. & Ward, J.D. 'Sausage toe': a reliable sign of underlying osteomyelitis, 74

Ramachandran, A. see Snehalatha, C.

Ramachandran, A. see Vijay, V.

Rani, G. see Deepa, R.

Ranstam, J. see Östgren, C.J.

Råstam, L. see Östgren, C.J.

Rayman, G. see Yuen, K.C.J.

Raymond, N.T. see Davies, M.J.

Reddy, A. see Yuen, K.C.J.

Reichard, P., Jensen-Urstad, K., Ericsson, M., Jensen-Urstad, M. & Lindblad, L.E. Autonomic neuropathy – a complication less pronounced in patients with Type 1 diabetes mellitus who have lower blood glucose levels, 860

Reichelt, A.J. see Schmidt, M.I.

Repetti, E. see Bo, S.

Reviewed By, N. & Finer, M.L. Type 1 diabetes mellitus and hepatic sinusoidal fibrosis, 89

Ricerca, B.M. see Ghirlanda, G.

Rich, S.S. see Moczulski, D.K.

Richard-Harston, S. see Bourdel-Marchasson, I.

Riche, N. see Munro, N.

Rigalleau, V., Gatta, B., Bonnaud, S., Masson, M., Bourgeois, M.L., Vergnot, V. & Gin, H. Diabetes as a result of atypical anti-psychotic drugs – a report of three cases, 484

Riis, A. see Home, P.D.

Roberts, A.B. see Cundy, T.:

Robinson, I. see Donohoe, M.E.

Robinson, R. see Morris, B.W.

Robinson, S. see Kousta, E.

Roep, B.P. see van Essen, E.H.R.

Rogers, C. see Morgan, C.Ll.

Rogus, J.J. see Moczulski, D.K.

Rohold, A. see Abrahamsen, B.

Rønningen, K.S. see Cinek, O.

Rooman, R.P.A. see De Block, C.E.M.

Rosén, I. see Sundkvist, G.

Rossing, K. see Clausen, P.

Rotella, C.M. see Mannucci, E.

Rowley, M.J. see Li, J.K.Y.

Royal College Of Physicians Of Edinburgh Diabetes Register Group. Near-normal urinary albumin concentrations predict progression to diabetic nephropathy in Type 1 diabetes mellitus, 782

Ruta, D.A. see Evans, J.M.M.

Rutten, G.E.H.M. see Van Den Arend, I.J.M.

Rutter, M.K., McComb, J.M., Forster, J., Brady, S. & Marshall, S.M. Increased left ventricular mass index and nocturnal systolic blood pressure in patients with Type 2 diabetes mellitus and microalbuminuria. 321

Ryan, A. see Nelson, M.

Ryder, R.E.J. see Kong, N.

Salmela, P.I. see Koivukangas, V.

Salo, S. see Tupola, S.

Sandbaek, A. see Meadows, K.A.

Santeusanio, F. see Buyken, A.E.

Sarioglu, N. see Kordonouri, O.

Satyavani, K. see Snehalatha, C.

Scaglione, L. see Bo, S.

Scanlon, P. see Griffin, M.E.

Schaper, F. see Hanefeld, M.

Schaupp, L. see Brunner, G.A.

Scheek, L.M. see Buter, H.

Scherstén, B. see Öestman, J.

Schmidt, M.I., Matos, M.C., Reichelt, A.J., Costa Forti, A., De Lima, L. & Duncan, B.B. For The Brazilian Gestational Diabetes Study Group. Prevalence of gestational diabetes mellitus – do the new WHO criteria make a difference?, 376

Schrijvers, G.J.P. see Van Den Arend, I.J.M.

Schultz, C.J., Dalton, R.N., Turner, C., Neil, H.A.W., Dunger, D. & For The Oxford Regional Prospective Study Group.

Freezing method affects the concentration and variability of urine proteins and the interpretation of data on microalbuminuria, 7

Schuppan, D. see Kordonouri, O.

Scott, A. Slide Atlas of Diabetes, 817

Scott, L. see Moczulski, D.K.

Seed, P.T. see Coster, S.

Seena, R. see Vijav, V.

Sehlin, J. see Oldenborg, P.-A.

Sendlhofer, G. see Brunner, G.A.

Shanthi, S. see Deepa, R.

Sharp, P.F. see Hipwell, J.H.

Shau, Y.-W. see Hsu, T.-C.

Shaw, K.M. see Meeking, D.R.

Shaw, N.J. see Ehtisham, S.

Shepherd, M. see Pearson, E.R.

Sherwood, R.A. see McIntyre, E.A.

Sima, A.A.F. see Sundkvist, G.

Simmons, D., Thompson, C.F. & Conroy, C. Incidence and risk factors for neonatal hypoglycaemia among women with gestational diabetes mellitus in South Auckland, 830

Singh, A.K., Bhatia, E., Dabadghao, P., Bhatia, V., Gellert, S.A. & Colman, P.G. Role of islet autoimmunity in the aetiology of different clinical subtypes of diabetes mellitus in young north Indians, 275

Singh, R.K. see Perros, P.

Sipilä, I. see Tupola, S.

Siu, S.T.S. see Janus, E.D.

Sivasankari, S. see Snehalatha, C.

Skyrme-Jones, R.A.P. & Meredith, I.T. Aspirin does not improve endothelial vasodilator function in Type 1 diabetes mellitus, 556

Smith, D. see McKenna, K.

Smith, J.M. see Whitelaw, D.C.

Smithers, M. see Morgan, C.Ll.

Snajderová, M. see Cinek, O. Snehalatha, C. see Vijay, V.

Snehalatha, C., Sivasankari, S., Satyavani, K., Vijay, V. & Ramachandran, A. Insulin resistance alone does not explain the clustering of cardiovascular risk factors in southern India,

Sochett, E.B. see O'Hayon, B.E.

Song, S.H., Mowbray, A., O'Shea, H. & Campbell, I.W. Bilateral optic atrophy following diabetic ketoacidosis, 394

Sönksen, P.H. see Coppini, D.V.

Sönksen, P.H. see Weng, C.

Sood, A. see Ahluwalia, A.

Sørensen, B. see Jensen, D.M.

Sowter, H. see Sturrock, N.D.C.

Spagnoli, A. see Chiarelli, F.

Spengler For The Finnish Acarbose Study Group, M. see Lindström, J.

Spoor, P.A. see Airey, C.M.

Squatrito, S. see Wolffenbuttel, B.H.R.

Standl, E. see Meier, M.

Stanton, K. see Collis, J.

Start, K. see Nelson, M.

Stead, J.W. see Donohoe, M.E.

Stehle, P. see Buyken, A.E.

Steinke, D. see Evans, J.

Stenhouse, E. see Zhao, H.X.

Stenhouse, E., Zaho, H.X. & Millward, B.A. Birthweight is not a risk factor for childhood-onset Type 1 diabetes mellitus, 552

Stevenson, J. see Sturrock, N.D.C.

Stewart, L.C. see McTernan, C.L.

Stolk, R.P. see Van Den Arend, I.J.M.

Storti, S. see Ghirlanda, G.

Stott, N.C.H. see Morgan, C.Ll.

Strachan, F. see Hipwell, J.H.

Strachan, M.W.J., MacCuish, A.C. & Frier, B.M. The care of students with insulin-treated diabetes mellitus living in university accommodation: scope for improvement?, 70

Stronge, J. see Griffin, M.E.

Sturrock, N.D.C., George, E., Pound, N., Stevenson, J., Peck, G.M. & Sowter, H. Non-dipping circadian blood pressure and renal impairment are associated with increased mortality in diabetes mellitus, 360

Suematsu, C. see Okada, K.

Sullivan, K. see Sundkvist, G.

Šumník, Z. see Cinek, O.

Sundkvist, G. see Öestman, J.

Sundkvist, G., Dahlin, L.-B., Nilsson, H., Eriksson, K.-F., Lindgärde, F., Rosén, I., Lattimer, S.A., Sima, A.A.F., Sullivan, K. & Greene, D.A. Sorbitol and *myo*-inositol levels and morphology of sural nerve in relation to peripheral nerve function and clinical neuropathy in men with diabetic, impaired, and normal glucose tolerance, 259

Sundqvist, I.-M. see Oldenborg, P.-A.

Sundsfjord, J. see Jorde, R.

Sutton, M. see Rajbhandari, S.

Suurinkeroinen, L. see Klannemark, M.

Swaminathan, R. see Coster, S.

Tai, E.S., Lim, S.C., Tan, B.Y., Chew, S.K., Heng, D. & Tan, C.E. Screening for diabetes mellitus – a two-step approach in individuals with impaired fasting glucose improves detection of those at risk of complications, 771

Takamura, T. see Nagai, Y.

Tan, B.Y. see Tai, E.S.

Tan, C.E. see Tai, E.S.

Tanaka, S. see Oki, Y.

Tang, F.-T. see Hsu, T.-C.

Taskinen, M.R. see Klannemark, M.

Taylor, R. Diabetic Retinopathy: Diagnosis, Management and Reference Images, 689

Taylor, R. see Collis, J.

Taylor, R. see Donohoe, M.E.

Taylor, R. see Pandit, R.J.

Temelkova-Kurktschiev, T. see Hanefeld, M.

Tesfaye, S. see Rajbhandari, S.

The Belgian Diabetes Registry, L.F. see De Block, C.E.M.

The Data Committee For Childhood Diabetes Of The Japan Diabetes Society (JDS), Z. see Kida, K.

Thomas, A. Standardization of HbA<sub>1c</sub> measurements - the issues,

Thomas, A., Middle, J., John, G., Marshall, S.M. & Barth, J.H. Standardization of HbA<sub>1c</sub> measurements. Reply, 559

Thomas, B.J. see Nelson, M.

Thompson, C.F. see Simmons, D.

Thompson, C.I. see McKenna, K.

Tildesley, H. see Meneilly, G.S.

Todaro, L. see Ghirlanda, G.

Toeller, M. see Buyken, A.E.

Tomkin, G.H. see Owens, D.

Tooke, J.E. see Donohoe, M.E.

Tormey, W. see McKenna, K.

Törn, C. see Öestman, J.

Townend, K. see Cundy, T.

Tsumura, K. see Okada, K.

Tumini, S. see Chiarelli, F.

Tuomilehto, J. see Lindström, J.

Tupola, S., Sipilä, I., Huttunen, N.P., Salo, S., Nuuja, A. & Åkerblom, H.K. Management of asymptomatic hypoglycaemia in children and adolescents with Type 1 diabetes mellitus,

Turner, B.C. see Browne, D.L.

Turner, C. see Schultz, C.J.

UK Prospective Diabetes Study (UKPDS) Group. Effects of three months' diet after diagnosis of Type 2 diabetes on plasma lipids and lipoproteins (UKPDS 45), 518

Unwin, N. see Wareham, N.J.

Unwin, N.C. see Levitt, N.S.

Urbancic-Rovan, V. & Gubina, M. Bacteria in superificial diabetic foot ulcers, 814

Vaag, A. see Fenger, M.

Valk, G.D. see Franse, L.V.

Vambergue, A., Nuttens, M.C., Verier-Mine, O., Dognin, C., Cappoen, J.P. & Fontaine, P. Is mild gestational hyperglycaemia associated with maternal and neonatal complications? The Diagest Study, 203

van den Arend, I.J.M., Stolk, R.P., Rutten, G.E.H.M. & Schrijvers, G.J.P. Education integrated into structured general practice care for Type 2 diabetic patients results in sustained improvement of disease knowledge and self-care, 190

van den Ouweland, J.M.W. see van Essen, E.H.R.

Van Eijk, J.T.M. see Franse, L.V.

van Essen, E.H.R., Roep, B.P., Hart, L.M.'t, Jansen, J.J., van den Ouweland, J.M.W., Lemkes, H.H.P.J. & Maassen, J.A. HLA-DQ polymorphism and degree of heteroplasmy of the A3243G mitochondrial DNA mutation in maternally inherited diabetes and deafness, 841

Van Gaal, L.F. see De Block, C.E.M.

Van Tol, A. see Buter, H.

Vaughan, N.J.A. & Fogarty, L. Confidentiality and diabetes 'registers' – a dilemma?, 563

Vavrinec, J. see Cinek, O.

Vergnot, V. see Rigalleau, V.

Verier-Mine, O. see Vambergue, A.

Verrotti, A. see Chiarelli, F.

Vijay, V. see Snehalatha, C.

Vijay, V., Narasimham, D.V.L., Seena, R., Snehalatha, C. & Ramachandran, A. Clinical profile of diabetic foot infections in south India - a retrospective study, 215

Wahren, J. see Johansson, B.-L.

Wake, M., Hesketh, K. & Cameron, F. The Child Health Questionnaire in children with diabetes: cross-sectional survey of parent and adolescent-reported functional health status,

Waldron, S. see Nelson, M.

Walker, D. see Khawaja, K.I.

Walker, L. see Nelson, M.

Walker, M. see New, J.P.

Wallymahmed, M.E. see Ismail, A.A.

Wang, C.-L. see Hsu, T.-C.

Wang, C.L. see Petry, C.J.

Ward, J.D. see Rajbhandari, S.

Wareham, N.I., Unwin, N. & Borch-Johnsen, K. Descriptive epidemiology short reports in Diabetic Medicine - an opportunity to present data with the population as the unit of variation, 691

Warram, J.H. see Moczulski, D.K.

Wat, N.M.S. see Janus, E.D.

Watkins, P.J. see Winkler, A.S.

Watkins, P.J. see Winkler, A.S.

Watson, J. see Flanagan, D.E.H.

Watts, G.F. see Dogra, G.

Weber, G. see Kordonouri, O.

Weiner, K. Potential for error from underfilling Glucometer Elite test strips, 555

Weng, C. see Coppini, D.V.

Weng, C., Coppini, D.V. & Sönksen, P.H. Geographic and social factors are related to increased morbidity and mortality rates in diabetic patients, 612

West, I.C. Radicals and oxidative stress in diabetes, 171

Westergaard, J.G. see Jensen, D.M.

Whitelaw, D.C., Clark, P.M., Smith, J.M. & Nattrass, M. Effects of the new oral hypoglycaemic agent nateglinide on insulin secretion in Type 2 diabetes mellitus, 225

Wibell, L. see Öestman, J.

Williams, B. see Meier, M.

Williams, D.D.R. see Airey, C.M.

Williams, R. see McKinney, P.A.

Winkler, A.S. & Watkins, P.J. Long term treatment of the anaemia in Type 1 diabetes mellitus with erythropoietin, 250

Winkler, A.S., Ejskjaer, N., Edmonds, M. & Watkins, P.J. Dissociated sensory loss in diabetic autonomic neuropathy,

Winnock, F. see De Block, C.E.M.

Wolffenbuttel, B.H.R., Gomis, R., Squatrito, S., Jones, N.P. & Patwardhan, R.N. Addition of low-dose rosiglitazone to sulphonylurea therapy improves glycaemic control in Type 2 diabetic patients, 40

Wu, Y .- C. see Jap, T .- S.

Wutte, A. see Brunner, G.A.

Yamashita, H. see Nagai, Y.

Yeong, M.L. see McGrath, N.M.

Young, P.J. see Coppini, D.V.

Yuen, K.C.J., Baker, N.R., Reddy, A., Edelsten, C. & Rayman, G. Blindness following a diabetic foot infection: a variant to the 'eye-foot syndrome'?, 546

Zaho, H.X. see Stenhouse, E.

Zhao, H.X., Stenhouse, E., Demaine, A.G. & Millward, B.A. The average age at onset is decreasing in children with Type 1 diabetes in Devon and Cornwall, England, 877

Zimmet, P. see Davis, T.M.E.

Zimmet, P.Z. see Li, J.K.Y.

Zouvanis, M. see Panz, V.R.

## Subject Index

17-β-estradiol, 308

ADA, 771 fasting criteria, 872 acarbose, 20 acetylcholine, 243

adhesion molecules, 644 adrenaline, 848 adrenoceptors, 848

adverse maternal and fetal outcome, 203 aetiology of Type 1 diabetes mellitus, 59 African subjects, 524

age at onset of diabetes, 538 aged, 346 albumin excretion rate, 708 alcohol, 70, 776 aldose reductase gene, 111 alpha-glucosidase, 810 ambulatory blood pressure, 321 monitoring, 360

American Diabetes Association, 381 amputations, 727

angiotensin converting enzyme inhibition, 134

angiotensin-converting enzyme inhibitors, animal insulin, 416

anxiety, 198 Arabs, 720

ascorbic acid, 171

Asians, 720 atherosclerosis, 299

atrial natriuretic peptide, 219, 463, 512 attitudes, 581

atypical anti-psychotic, 484 audit, 565

autoimmunity, 841 autonomic, 163 nervous system, 606 neuropathy, 457, 860

B-cell function, 445 bacteraemia, 546 beta cell dysfunction, 433 biomechanics, 854 birth length, 158 birth weight, 158 birthweight, 48, 365, 830 blindness, 727 blood glucose self-monitoring, 755 blood glucose, 269 blood pressure, 606 blood pressure, 782 body composition, 124 body mass index, 48, 478, 675 bone mineral density, 124 brain natriuretic peptide, 512

CVD risk factors, 152 capillary, 163 carbohydrate intake, 351 carbon monoxide, 119 cardiovascular, 644 disease, 124 risk factors, 835 risk, 741 carotid ultrasound, 299 cell cholesterol, 308 cerebral oedema, 807 Charcot foot, 253 chemiluminescence, 532 child, 700 childhood, 650 Type 1 diabetes mellitus, 59, 236 children, 867 Chinese, 295, 798 cholesterol, 518 ester transfer protein, 308 cigarette smoking, 119 clinical guidelines, 565 clinical standards, 565 clozapine, 484 coeliac disease, 441, 657 community pharmacists, 528 compliance, 528 complications, 146 of diabetes, 253 coping, 636 cost analysis, 627 cotinine, 119 C-peptide, 269 cross-cultural adaptation, 572 cross-sectional study, 776

cystoid macular oedema, 326

Czech, 64

DPP-IV. 346 day care, 236 'dead in bed' syndrome, 70 decision analysis, 469 decision-making, 823 depression, 198 diabetes education, 727 diabetes in relatives other than parents, diabetes mellitus, 15,74, 130, 171, 219, 386, 416, 451, 478, 481, 495, 512, 572, 810,848 drug therapy, 675 subtypes, 524 diabetes, 119, 141, 146, 163, 230, 441, 484, 532, 581, 700, 741 outcome, 612 screening, 771 related psychological distress and behadiabetic coma, 70 diabetic foot, 215, 854 diabetic ketoacidosis, 295, 394, 807 diabetic nephropathy, 111, 644, 650, 708 diabetic neuropathies, 105 diabetic retinopathy, 627 diagnostic criteria, 381, 741 diastolic function, 321 diazoxide, 91 diet, 518 dietary carbohydrates, 20, 336 dietary fat, 20 digital fundus photography, 627 digital retinal photography, 588 dipping, 360 driving, 316 dyslipidaemia, 599

education, 581, 636, 682 elderly, 675 emergency service, 386 endogenous staphylococcal endophthalmitis, 546 endomysial antibodies, 441 endothelium, 243, 644 environmental, 612 epidemiology, 776 ethnicity, 667, 792 euglycaemic hyperinsulinaemic clamp, 713 evidence-based medicine, 823 exercise, 53 eye-foot syndrome, 546

fMet-Leu-Phe, 532 factor analysis, 152 familiar diabetes, 538 fasting glucose, 381, 771 fasting hyperglycaemia, 433 fasting plasma glucose, 287, 595 female–male ratio, 59 fetal outcome, 26 fibrinogen, 798 fibro-calculous pancreatic diabetes, 275 food groups, 351 foot complications, 215, 215 forearm blood flow, 243

GAD antibody, 275

GAD autoantibodies, 524 GADA, 618 GIP, 346 GLP-1, 346 general practice, 230 genetic susceptability, 111 genetics, 543, 735, 841 gestational diabetes mellitus, 26, 33, 281, gestational diabetes, 720, 792, 830 gestational impaired glucose tolerance, 48 glaucoma, 693 glomerular filtration rate, 181, 708 glucagon-like peptide-1, 713 glucocorticoids, 15 glucose, 720 metabolism, 124 tolerance test, 381 glutamic acid decarboxylase, 667 glutathione, 171 gluten-free foods, 657 glycaemic control, 40, 198, 636, 650, 782 glycaemic index, 336, 657 glycogen storage disease type III, 810

HbA<sub>1c</sub>, 351, 451 HLA DQ haplotype, 618 HLA, 661 HLA, 814 HNF-1α, 543 HOMA, 299 hand neuropathy, 682 health education, 823 health status, 700 heart rate variability, 860 heart transplantation, 15 heel, 854 hepatocyte nuclear factor-1α, 390 Hong Kong, 798 hormone replacement therapy, 308 human diabetes, 259 human insulin, 416 human muscle glycogen synthase, 735 hyperinsulinaemia, 365, 606 hyperinsulinaemism, 91 hypertension, 134, 152, 798 hypoglycaemia, 70, 209, 316, 416, 762

IA-2 antibody, 275 ICA512/IA-2, 667



image processing, 588 impaired glucose tolerance, 365, 433, 776 incidence, 59, 64, 675 infected foot ulcer, 546 infections, 236 inhibitor, 810 injection-meal interval, 371 insulin, 124, 316, 336, 433 analogue, 371, 762 analogues, 209 aspart, 371, 762 injection regime, 351 resistance, 152, 299, 433, 735, 867 secretion, 91, 225 sensitivity, 134 therapy, 209, 371, 762 integrated footcare, 581 intensified treatment, 860 intima-media, 835 intraocular pressure, 693 ischaemia, 394 isolated post-challenge hyperglycaemia, 835

joint clinic, 481

K<sub>ATP</sub> channels, 91 ketoacidosis, 326 knowledge, 528, 581

LDL fatty acids, 308
LDL-cholesterol, 538
LPL291, 599
laser-Doppler flowmetry, 130
left ventricular hypertrophy, 321
leisure activities, 53
leptin, 713
liaison psychiatry, 481
lifestyle, 53
lipoate, 171
lipoprotein composition, 308
lipoprotein lipase, 599
liver dysfunction, 810
longitudinal study, 451
low protein diet, 848

MODY, 553
MODY3 gene, 390
macrovascular, 146
male predominance, 64
malnutrition-modulated diabetes mellitus, 275
maternal age, 158
maternally inherited diabetes and deafness, 841
meta-analysis, 755
metabolic control, 20
metabolic syndrome, 365
metformin, 507
microalbuminuria, 7, 181, 219, 321, 463, 644, 782

microaneurysms, 588
microcirculation, 130
microvascular, 146
mild gestational hyperglycaemia, 203
mitochondria, 841
morbidity, 612
mortality, 612, 727
motivation, 636
mydriasis, 693

nateglinide, 225
neonatal hypoglycaemia, 91, 830
neonatal morbitity, 507
nephromegaly, 708
nephropathy, 727, 360
nerve fibre, 163
neurological examination, 105
neuropathy, 163, 215, 253, 394
non-immunological contact urticaria, 130
noradrenaline, 848
norethisterone acetate, 308

OGTT, 433, 771 obesity, 346, 478, 713, 798, 867 offspring, 606 olanzapine, 484 optic disc, 394 oral glucose tolerance test, 203, 595, 741 oral hypoglycaemic agents, 281, 507, 528 osteomyelitis, 74 oxidative stress, 171

PVD, 215 Palestinian population, 746 pancreatic function, 295 parity, 158 patient education, 190 perinatal mortality, 33 peripheral nerve function, 259 peroxidase, 532 peroxisomal proliferator-activated receptor-y (PPARy), 287 pharmacogenetics, 553 pilocarpine, 693 placental ratio, 48 placental weight, 48 plasma cholesterol, 308 plethysmography, 243 polymorphonuclear leucocytes, 532 Polynesians, 830 population-based survey, 675 post-prandial hyperglycaemia, 835 postural hypotension, 163 pre-eclampsia, 507 pregnancy, 26, 33, 281, 507, 830 prevalence, 7, 26, 141, 376, 478, 746 previous gestational diabetes mellitus, 595 primary care, 190 primary healthcare, 105

programme planning, 627

prospective studies, 53 proteinuria, 360 psychosocial, 700 support, 636

QALYs, 469 quality of care, 565 quality of life, 700 quality of life, 762

randomized controlled trial, 209 rapid-acting insulin, 762 reactive oxygen species, 171 record linkage, 141, 146 reference values, 7 referrals, 230 reflex sympathetic dystrophy, 253 relative activity, 230 reliability, 572 respiratory heart rate variability, 181 retinopathy, 495, 782 risk communication, 823 risk factors, 376, 746 rosiglitazone, 40, 287

S-100B protein, 807 sausage toe, 74, 469, 495, 627, 720 seasonality, 59 self-care behaviour, 190 short stature, 792 small fibre polyneuropathy, 457 smoking, 445, 727 social deprivation, 612 socioeconomic factors, 478 socioeconomic status, 727 soluble human insulin, 371 sorbitol, 259 south Indians, 872 spectral analysis, 606 structured care, 190 substance P, 243 sulphonylurea, 40, 91, 507 sensitivity, 553 superoxide, 532 sural nerve, 259 survival, 360 sympathectomy, 253 syndrome 'X', 152 systematic review, 495, 416, 755

Taiwan, 390 telephone helpline, 386 tertiary education, 70 thermal thresholds, 181 thiazolidinedione, 40, 287 thyrogastric antibodies, 618 tissue transglutamase antibodies, 441 triglyceride, 518, 693 twin-control study, 158 twins, 158, 365, 735

Type 1 diabetes mellitus, 7, 64, 70, 111, 158, 198, 209, 269, 275, 351, 394, 457, 463, 661, 667, 762, 782, 841, 860

Type 2 diabetes mellitus, 20, 33, 40, 105, 119, 134, 190, 198, 225, 269, 287, 308, 321, 346, 445, 469, 518, 528, 538, 546, 599, 667, 713, 735, 746, 755, 776, 798, 823, 830, 841, 867

UKPDS, 518, 823 ulceration, 682, 854 ultrasonography, 854 universal screening, 26 uveitis, 326

validity, 572 variability, 7 vascular endothelial growth factor, 650 vascular reactivity, 130 vitamin E, 171 vitreous fluorophotometry, 181 voglibose, 810 WHO, 771 2-h criteria, 872 World Health Organization, 381

